

Histological study about *Leishmania infantum* infection on canine inflammatory bowel disease

UAB

Universitat Autònoma
de Barcelona

Sandra Martín Nicolau – June 2016




FACULTAT DE VETERINÀRIA

INTRODUCTION

The inflammatory bowel disease (IBD) is an idiopathic inflammatory disorder that affects gastrointestinal tract (GIT). The histopathological evaluation shows an inflammation in the colonic mucosa: lymphoplasmacytic colitis and histiocytic colitis. A pathogen or infectious agent could be a possible cause of IBD such as *Leishmania infantum*. The **main aim** of this study was to clarify if *L. infantum* infection would be a possible cause of IBD.

MATERIAL AND METHODS

1. Database selection



2. Immunohistochemistry for *Leishmania*



SPECIFIC OBJECTIVES AND RESULTS

1. To retrospectively evaluate the diagnoses of *L. infantum* infection in dogs with IBD.




Figure 1. Percentages of *Leishmania* IHC performed in dogs diagnosed with IBD and percentages of *Leishmania* IHC results from Diagnet.

Inflammation pattern	Percentage
Lymphoplasmacytic enteritis	32.6 %
Lymphoplasmacytic colitis	19.4 %
Histiocytic colitis	2.6 %
Histiocytic enteritis	0.4 %

Table 1. Percentage of types of inflammation and the segment of the gastrointestinal tract affected in dogs with a diagnosis of IBD from Diagnet.

2. To determine *L. infantum* infection in colonic biopsies diagnosed with IBD and description of the clinical cases.

Detection of 1 positive case from 27 cases by IHQ technique (3.6 %).

Case	Breed	Sex	Age (years)	Presenting complaint	Endoscopic findings	Other observations
Dog 1	German Dogo	M	9	Large bowel diarrhoea	↑ Nematodes	Diagnosis of <i>Leishmania</i> 3 months before biopsy
Dog 2	Boxer	M	5	Chronic diarrhoea with hematochezia	-	Echography: mesenteric iliac lymphadenomegaly
Dog 3	Cocker Spaniel	M	15	Chronic hematochezia	-	-
Dog 4	Mixed breed	F	1	Chronic diarrhea with mucus and bleeding for 7 months. Anorexia for 3 days, apathy and weakness	Hypertrophy, hyperemia and mucosal petechial	Echography: alteration of duodenum and colon
Dog 5 (new +)	Belgian Shepherd	F c	12	Bloody diarrhea for 1 month	Duodenum and colon very altered with a friable and edematous mucosa	Diagnosed of <i>Leishmania</i> before biopsy by the clinical veterinarian

Table 2. Description of the clinical cases positive to *Leishmania* by IHC.

3. Description of the histological lesions and parasite load of cases diagnosed with IBD and infection with *Leishmania*.




Figure 2. Positive case. Histological sections of the colon. A: intense histiocytic inflammation, HE staining. B: the same case than in the image A, but with IHC. Intense parasitization, amastigotes can be seen throughout the image. Arrows indicate some of amastigotes of *Leishmania* that can be seen.

Case	Year	Inflammation	Grade of intensity of inflammation	Amastigote count (counting 5 fields 40x)
Dog 1	1997	lymphoplasmacytic	Intense	33
Dog 2	2004	lymphoplasmacytic	Mild	37
Dog 3	2011	Histiocytic	Intense	15
Dog 4	2013	Histiocytic	Intense	Not able to count due to massive number of amastigotes
Dog 5 (new +)	2007	lymphoplasmacytic	Mild	2

Table 3. Description of IHC positive cases by *Leishmania*.

CONCLUSIONS

- The inflammation associated with *Leishmania* infection in the colon is both histiocytic and lymphoplasmacytic.
- Both the degree of inflammation and parasitism seems to vary between individuals.
- IHC for *Leishmania* is not commonly employed to describe this infection in dogs with a histological diagnosis of IBD.
- IHC for *Leishmania* should be routinely used as a diagnostic tool in endemic areas, to exclude or confirm an infection by this parasite in patients with a diagnosis of IBD.